

Hanging Wet-to-Dry Dressings Out to Dry

Advances in Skin & Wound Care: The Journal for

Liza G Ovington

Prevention and Healing

March/April 2002

Volume 15 Number 2

Pages 79 - 84

PURPOSE To present information on effective alternatives to wet-to-dry dressings in the management of wounds.

TARGET AUDIENCE This CME/CE activity is intended for physicians and nurses with an interest in alternatives to wet-to-dry dressings in the management of wounds.

OBJECTIVES

Identify factors that affect wound healing.

- 1.
2. Discuss advantages and disadvantages of various types of wound care products.
3. Describe strategies to promote cost-effective wound healing.

Health care professionals managing patients with chronic or complex wounds have a wide and exciting variety of treatment options and adjunctive therapies from which to choose. The list of products that have shown positive benefits for wound healing includes a recombinant protein drug for diabetic foot ulcers, bioengineered tissue products for venous and diabetic ulcers, negative pressure therapy, normothermic therapy, constant tension approximation, electrical stimulation, hyperbaric oxygen, and hundreds of topical dressings.

Wound dressings in particular have undergone an extensive diversification over the past quarter century. In the 1960s, animal and human studies showed that wounds in which the tissues remained moist healed twice as rapidly as those in which the tissues dried out. ¹⁻³ Soon manufacturers embraced this new moist wound healing concept and began to develop dressings from materials that could prevent wound tissues from desiccating. Gauze dressings have been joined, but not displaced, by polymeric materials such as polyurethane films, foams, hydrogels, and hydrocolloids, as well as calcium alginates, collagens, and other materials. These dressings are often collectively referred to as "moisture-retentive" or "semi-occlusive." Today, over 50 manufacturers produce more than 350 brands of moisture-retentive wound dressings.

With so many choices for topical wound management, one might expect to find a wide variety of dressing products in use among wound patients at any one facility or in any particular care setting. However, this is not the norm. Despite the benefits of the newer dressing products, gauze is still the most widely used wound care dressing and may be erroneously considered a standard of care.

In an illustrative study, 13 home care agencies in 1 geographic area gathered data regarding the topical dressings being used during 1 week for 1029 patients with 1638 classified wounds. -Wound types included chronic wounds such as pressure ulcers, venous leg ulcers, and diabetic foot ulcers as well as surgical wounds and other skin tears. The survey found that the most commonly used dressing for all types of wounds was gauze—dry gauze. The majority of cases (n = 406) were treated with dry gauze.

Saline-moistened gauze was the third most-used dressing (n = 145), with the second most common being no dressing at all (n = 252). Advanced moisture-retentive dressings such as alginates, collagens, films, foams, hydrogels, and hydrocolloids accounted for less than 25% of all dressings used.

One could argue that dry gauze dressings have little value for optimal wound management in today's health care environment. Even saline-moistened gauze dressings have little value when compared with the more advanced dressing categories. Gauze dressings (whether dry or moistened with saline) are substandard for optimal wound care for several reasons:

1. they may actually add to the patient's discomfort, impede healing, and increase the risk of infection;
2. they may incur more labor for the clinician or caregiver; and
3. they may incur more costs for the health care system.

This article examines each of these allegations in detail.

Wet-to-Dry and Wet-to-Moist: What's the Difference?

Wet-to-dry and *wet-to-moist* are terms used in conjunction with gauze dressings and technically they are two distinctly different ways of using saline (the most common wetting agent). However, in actual use, there may be little difference between the two.

Wet-to-dry gauze dressings

Wet-to-dry gauze dressings are typically intended for use in the debridement of devitalized tissues from the wound bed. Gauze is moistened with normal saline and placed into or onto the wound. This moistened layer is then covered with dry layers of gauze or other dry dressings (such as abdominal pads). The moistened gauze then dries out; while drying, it adheres to surface tissues, which are removed when the dried dressing is removed.

Wet-to-dry debridement is not selective and often also removes healthy tissues, causing reinjury. The patient may also suffer significant pain upon removal of a wet-to-dry dressing. Alternative forms of debridement such as surgical, sharp, enzymatic, and autolytic offer a higher degree of selectivity and less pain in most cases. *Wet-to-moist gauze dressing*

A wet-to-moist dressing is generally prepared in the same manner as a wet-to-dry dressing; however, it is intended to remain continuously moist until removal. Nevertheless, it may become a wet-to-dry dressing in practice. A recent study of the mechanism of action of saline dressings suggests that they function as an osmotic dressing.⁵ Normal saline is isotonic. As water evaporates from the saline dressing, it becomes hypertonic and fluid from the wound tissues is drawn into the dressing in an attempt to reestablish isotonicity.

However, wound fluid is not merely water; it contains blood and proteins that may begin to form an impermeable layer on the dressing's surface. At this point, fluid from the wound is unable to replace the fluid lost from the dressing by evaporation and the dressing dries out completely. Therefore, unless careful attention is paid to the moisture levels in a wet-to-moist gauze dressing—either through frequent changing of the dressing or through remoistening of the gauze with additional saline—it becomes wet-to-dry.

Removal of a wet-to-moist dressing that has dried may then cause reinjury of the wound, resulting in pain and delayed wound healing. Semiocclusive dressings that maintain a moist wound environment have been associated with less pain at dressing change than gauze dressings, as well as less pain in general. - '-The diminution of pain

is based on the fact that nerve endings in the wound tissue are protected from exposure and dehydration.

Gauze Dressings: Patient Issues

Impeded healing—local tissue cooling

Another effect of a gauze dressing drying out is that of local tissue cooling. In general, evaporation of water from a surface results in a reduction of temperature at that surface due to water's high heat of vaporization (540 cal per gram). In an open wound with nothing to impede fluid evaporation, the tissue temperature has been measured at 21°C. A gauze dressing placed in the wound does little to impede fluid evaporation and tissue temperature measures 25°C to 27°C—still approximately 10° below normal tissue temperature. -

The use of a semioclusive dressing, which impedes moisture loss from the wound, will diminish this local cooling. Wounds dressed with transparent films or foams, for example, have tissue temperatures measuring 33°C to 35°C. Reductions in tissue temperature have multiple physiologic effects, including local reflex vasoconstriction and hypoxia, impairment of leukocyte mobility and phagocytic efficiency, and increased affinity of hemoglobin for oxygen—all of which not only impede healing but increase susceptibility to infection. A gauze dressing does not support healing; a dressing that reduces loss of the wound temperature facilitates healing.

Increased risk of infection

Gauze dressings present no physical barrier to the entry of exogenous bacteria. One dramatic in vitro study showed that bacteria were capable of penetrating up to 64 layers of dry gauze.² Moistened gauze presents even less of a barrier to bacterial penetration. This is an issue as more wounds are managed in the home environment where infection control is not rigorously practiced. This lack of infection control is further complicated by reports from the Centers for Disease Control that 40% of the population sheds *Staphylococcus aureus* from the nares.^{1,-}

Therefore, patients whose wounds are dressed with gauze dressings would appear to be more susceptible to wound infection than those whose wounds are dressed with a moisture-retentive dressing. Indeed, both retrospective and prospective clinical studies have shown that infection rates in wounds dressed with gauze are actually higher than in wounds dressed with transparent films or hydrocolloids. —'— For example, in a literature review of 3047 wounds, the overall infection rate for wounds dressed with moisture-retentive dressings was 2.6%, whereas the infection rate for gauze-dressed wounds was 7.1%.¹³

The rationale for this difference is thought to be the barrier effects of moisture-retentive dressings, as well as their ability to preserve the viability and function of endogenous phagocytes by maintaining a moist environment. —Although moisture-retentive dressings allow the passage of moisture vapor through their surfaces, they are impervious to liquids. Therefore, a moisture-retentive dressing with an adhesive border or secondary dressing will allow the patient to bathe or shower without disturbing the wound.

Gauze Dressings: Clinician and Caregiver Issues

Labor intensive

As explained previously, in order for gauze to remain continuously moist to support optimal healing, it must either be changed frequently or remoistened with additional saline. This requires additional labor on the part of the clinician or the lay caregiver.

Dressing changes 2 or 3 times a day used to be common. In today's reimbursement climate, this practice is no longer feasible, not only from a reimbursement perspective, but also from the standpoint of best patient outcomes.

In home health care, multiple dressing changes a day require the expense of extra travel, in-home visits, and post-visit time for documentation. These are not considerations in acute or long-term care but in those settings, frequent dressing changes still require time that could be used for other patient care tasks. In addition, frequent dressing changes have been associated with evaporative cooling of the wound as it is exposed to the air.

Distribution of airborne bacteria leading to cross contamination

The removal of a dried dressing from a wound disperses significant amounts of bacteria into the air. One study used hand-held air samplers to quantify the number of bacteria released from colonized wounds upon removal of gauze dressings from burns compared to the removal of hydrocolloid dressings.¹⁵ The airborne dispersal of bacteria was greatest when removing dry gauze dressings; however, even removing moist gauze dressings released bacteria. Removal of a moisture-retentive hydrocolloid dressing released almost no bacteria. The decline of these airborne bacteria was slow, taking up to 30 minutes.

Most chronic wounds have polymicrobial colonization. — '—Implications for potential cross-infection are considerable, especially if the patient has multiple wounds or if multiple patients are seen in the same area, simultaneously or consecutively. If the wound is colonized with antibiotic-resistant bacteria, the implications may be even more serious.

Gauze Dressings: Health Care System Issues

Because prospective payment has become the reality in most health care facilities and systems, everyone is looking for ways to decrease costs. Supply costs are a tangible and attractive target for cost savings and it may be tempting to think that a product that costs less will decrease costs of care. Those still using gauze dressings may feel financially impeded to move toward using advanced dressings, and even enlightened clinicians may feel that they can no longer afford to purchase advanced dressings. This thinking is wrong. Now more than ever, advanced dressings are financially sensible in wound care. Although dressings such as films, foams, hydrocolloids, hydrogels, collagen, and alginates, may be more expensive per dressing than gauze dressings, they are less expensive to use. It is important not to confuse the price of a product with the cost of care.

The real cost of wound care can be considered as the sum of:

- the price of the dressing
- the labor cost of having a health care professional change the dressing
- the indirect costs of ancillary supplies and services used in changing the dressing (eg, gloves and biohazardous waste disposal)
- the cost of duration of care (eg, facility charges and travel costs for home care).

Several studies have been done to establish that advanced wound dressings can be cost-effective simply by taking into account the cost of labor. A more expensive dressing that requires less frequent dressing changes and results in shorter healing times has been found to be much less expensive to use.^{1,9}

Colwell et al "demonstrated that even when the cost of the semioclusive dressing and ancillary supplies was \$6.15 per dressing change versus \$0.47 for wet-to-moist gauze,

the daily cost of care for the semiocclusive dressing was only \$3.55 versus \$12.26 for the gauze because the former required less frequent changes than the latter. Xakellis and Chrischilles -^c showed that while the materials cost of a semiocclusive dressing was more than 3 times higher than saline gauze, the nursing time required for use was one-eighth that of the saline gauze. The cost in use of the semiocclusive dressing according to national nursing wages at the time was \$15.90 versus \$25.31 for the gauze dressing.

Bolton ²¹ proposed that a true, total cost of care should include yet another variable: the quality of the healing outcome in terms of clinical effect. Adding this variable underscores the idea that an inexpensive product is not really inexpensive if it does not produce the desired results (ie, timely healing and improved quality of life for the patient).

These researchers found moisture-retentive dressings actually decrease the costs of care relative to gauze, primarily through their impact on clinician labor, but also as a consequence of improved healing. Add to this the implications for decreased costs related to a lower incidence of infections and pain and semiocclusive dressings are a more logical alternative than ever before.

Illustrative Case Study

Table 1 is an example of the financial costs of managing a home health care patient with a wound. The table compares the use of gauze and saline for wound management versus more expensive advanced dressings that maintain a moist wound environment and better facilitate healing.

Table 1. COMPARISON OF COST AND EFFECTIVENESS OF 2 DRESSING TYPES IN A HOME HEALTH CARE PATIENT (TWICE-DAILY GAUZE DRESSING CHANGES)

Approximate costs for the supplies in this example were taken from the on-line catalog of a national medical product distributor. This estimate assumes a twice-daily dressing change for the saline and gauze dressing and a 3 times-per-week dressing change for the alternative—an adhesive-bordered foam dressing. Note that even for a dressing that is more than 10 times the price of gauze (\$10 versus \$0.75), the weekly materials cost is less than gauze due to the frequency of dressing changes. The table further shows that if the patient is occasionally doing dressing changes independently, the supply cost of gauze dressings, which the home health care agency would assume, is more than twice as much as the supply cost of an advanced dressing. Factor in the cost of the nursing labor at \$100 per visit to perform the dressing change. The labor costs for the advanced dressing are approximately 20% of the labor required for the gauze dressing changes. The weekly labor and materials costs for caring for the wound with gauze dressings is almost 5 times the costs of using the more expensive foam dressing. *Patient outcome*

Consider the cost-effectiveness of the 2 dressing modalities. Assume, based on the literature, that use of the advanced dressing results in more expedient healing of the wound. Assume that the wound is healed within 4 weeks of treatment with the advanced dressing and only 50% healed in the same amount of time with the gauze dressing. To determine cost-effectiveness, compute the cost of the effect of the treatment (ie, what did it cost in terms of labor and materials to attain a 1% reduction in the size of the wound?).

To obtain this cost figure, multiply weekly cost per treatment by 4 (for 4 weeks of treatments) and divide the total cost by the percent of healing achieved (either 50% or 100%). In this example, using gauze and saline to achieve healing costs more than \$115 per 1% reduction in wound size, whereas the semioclusive foam dressing costs approximately \$13 for the same reduction. As this typical example shows, far from saving money, gauze actually may be more expensive to use than a more effective dressing that costs 10 times its price. In [Table 2](#), the frequency of dressing changes for the saline-moistened gauze is daily instead of twice daily. There was still a significant cost difference in total care and rate of healing.

Table 2. COMPARISON OF COST AND

EFFECTIVENESS OF 2 DRESSING TYPES IN A HOME HEALTH CARE PATIENT (DAILY GAUZE DRESSING CHANGES

Clinicians who want to analyze the cost of caring for patients with wounds should put this table in a spread-sheet format to better anticipate the costs of wound care. As the plan of care is initially developed, the clinician should discuss the plans with the other members of the wound care team and patient and look prospectively and retrospectively at practices and costs.

Knowing All This—Now What?

Gauze dressings are not an optimal wound care modality for the patient, the clinician, or the health care system. They do not effectively support optimal healing and are more labor intensive to use than advanced dressings.

Reasons for the persistence of gauze as a wound management material may be related to several factors:

- Gauze and saline are essentially a one size fits all modality, are readily available and familiar, are perceived as inexpensive, and are the dressings of long tradition.
- Most advanced dressings are of discrete dimensions and cannot always be adjusted for wounds of different sizes, requiring that the health care facility stock multiple sizes.
- Not all physicians, nurses, and therapists have knowledge of the broad array of products available and understand the way they work.
- There are many brands and variations of advanced dressings that have variable appearances and performance, which may initially confuse the health care provider.
- As mentioned earlier, advanced dressings may also be incorrectly perceived as more expensive than gauze due to their individual purchase price.

Properly used, advanced dressings have significant benefits and should become the standard of care for wound management. Due to their diversity, the effective and efficient use of moisture-retentive dressings will require that clinicians to familiarize themselves with the range of dressing types and functions. Manufacturers provide specific information in product package inserts, instructions for use, and material safety data sheets. There are also multiple reviews of these dressings in the literature.

Why Gauze Dressings Should NOT Be Used:

- A gauze dressing placed in the wound does little to impede fluid evaporation and allows a loss of tissue temperature, resulting in impaired healing.
- Wet-to-dry debridement is not selective and often also removes healthy tissues, causing reinjury and significant pain.

Both retrospective and prospective clinical studies have shown that infection rates in wounds dressed with gauze are actually higher than in wounds dressed with transparent films or hydrocolloids.

In today's reimbursement climate, twice-daily and 3-times daily dressings are not feasible, not just from a reimbursement perspective, but also as an ineffective way to reach the most positive outcomes.

Removing a dried dressing from a wound disperses significant amounts of bacteria into the air. Research has determined that airborne dispersal of bacteria was greatest from dry gauze dressings; however, even removing moist gauze dressings released bacteria.

Semiocclusive dressings are more financially feasible from a total cost perspective.

References

1. Hinman C, Maibach H. Effect of air exposure and occlusion on experimental human skin wounds. *Nature* 1963;200:377.
2. Winter GD. Formation of the scab and the rate of epithelization of superficial wounds in the skin of the young domestic pig. *Nature* 1962;193:293.
3. Winter GD, Scales JT. Effect of air-drying and dressings on the surface of a wound. *Nature* 1963; 197:91.
4. Pieper B, Templin TN, Dobal M, Jacox A. Wound prevalence, types, and treatments in home care. *Adv Wound Care* 1999; 12:117-26.
5. Kim JK, Saliba L, Smith MJ. Normal saline wound dressing—is it really normal? *Br J Plast Surg* 2000;53:42-5.
6. Hedman LA. Effect of a hydrocolloid dressing on the pain level from abrasions on the feet during intensive marching. *Military Med* 1988; 153:188-90.
7. Nemeth AJ, Eaglstein WH, Taylor JR, Peerson U, Falanga V. Faster healing and less pain in skin biopsy sites treated with an occlusive dressing. *Arch Dermatol* 1991;117:1679-83.
8. Thomas S. *Wound management and dressings*. London, UK: The Pharmaceutical Press; 1990
9. Lawrence JC. Dressings and wound infection. *Am J Surg*; 1994;167:(Suppl 1A):21S-24S.
10. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practice Advisory Committee. *Am J Infect Control* 1999;27(2):97-132, (quiz 133-4, discussion 96).
11. Hutchinson JJ. Prevalence of wound infection under occlusive dressings, a collective survey of reported research. *Wounds* 1989(1):123-33.
12. Hutchinson JJ. A prospective clinical trial of wound dressings to investigate the rate of infection under occlusion. In: Harding K (editor). *Proceedings of the first European conference on advances in wound management*. London, UK: MacMillan; 1993. p 93-6.
13. Hutchinson JJ, McGuckin M. Occlusive dressings: a microbiologic and clinical review. *Am J Infect Control* 1990;18(4):257-68.
14. Mertz PM. Occlusive wound dressings to prevent bacterial invasion and wound infection. *J Am Acad Dermatol* 1985;12:662-8.
15. Lawrence JC, Lilly HA, Kidson A. Wound dressings and airborne dispersal of bacteria. *Lancet* 1992;339:807.
16. Brook I, Frazier EH. Aerobic and anaerobic microbiology of chronic venous ulcers. *Intl J Dermatol* 1998;37:426-8.
17. Bowler PG, Davies BJ. The microbiology of infected and noninfected leg ulcers. *Intl J Dermatol* 1999;38:573-8.
18. Alterescu V. The financial costs of inpatient pressure ulcers to an acute care

facility. *Decubitus* 1989;2(3): 14-23.

19. Colwell JC, Foreman MD, Trotter JP. A comparison of the efficacy and cost effectiveness of two methods of managing pressure ulcers. *Decubitus* 1993;6(4):28-36.
20. Xakellis GC, Chrischilles EA. Hydrocolloid versus saline gauze dressings in treating pressure ulcers: a cost effectiveness analysis. *Arch Phys Med Rehabil* 1992;73:463-9.
21. Bolton LL, van Rijswijk L, Shaffer FA. Quality wound care equals cost-effective wound care: a clinical model. *Adv Wound Care* 1997;10(4):33-8.

The author has disclosed that she is a consultant for and on the speaker's bureau of ETHICON, Inc, and is a stock shareholder of Johnson & Johnson, Inc. Reprinted with permission from Lippincott Williams & Wilkins: Ovington LG. Hanging wet-to-dry dressings out to dry. *Home Healthcare Nurse* 2001;19(8):477-84. *Adv Skin Wound Care* 2002 March/April; 15(2): 79-84 *Copyright © 2002 Lippincott Williams & Wilkins All rights reserved*